



# **BRIJUNI CONFERENCE**

**Space, Time and Life**

**26 - 30 August 20002**

## **BOOK OF ABSTRACTS**

**Editor: S. Danko Bosanac**

**DAVID ADVANCED RESEARCH WORKSHOP**

## REPORT DOCUMENTATION PAGE

Form Approved OMB No. 0704-0188

maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.

<b>1. REPORT DATE (DD-MM-YYYY)</b> 16-12-2002		<b>2. REPORT TYPE</b> Conference Book of Abstracts		<b>3. DATES COVERED (From - To)</b> 26 August 2002 - 30 August 2002	
<b>4. TITLE AND SUBTITLE</b>  Space, Time And Life				<b>5a. CONTRACT NUMBER</b> F61775-01-WF037	
				<b>5b. GRANT NUMBER</b>	
				<b>5c. PROGRAM ELEMENT NUMBER</b>	
<b>6. AUTHOR(S)</b>  S. D. Bosanac, editor				<b>5d. PROJECT NUMBER</b>	
				<b>5d. TASK NUMBER</b>	
				<b>5e. WORK UNIT NUMBER</b>	
<b>7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)</b> R. Boskovic Institute Bijenicka c. 54 Zagreb 10001 Croatia				<b>8. PERFORMING ORGANIZATION REPORT NUMBER</b>  N/A	
<b>9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)</b>  EOARD PSC 802 BOX 14 FPO 09499-0014				<b>10. SPONSOR/MONITOR'S ACRONYM(S)</b>	
				<b>11. SPONSOR/MONITOR'S REPORT NUMBER(S)</b> CSP 01-5037	
<b>12. DISTRIBUTION/AVAILABILITY STATEMENT</b>  Approved for public release; distribution is unlimited.					
<b>13. SUPPLEMENTARY NOTES</b>					
<b>14. ABSTRACT</b>  The Book of Abstracts for papers presented: Space, time and life, 26 August 2002 - 30 August 2002  Structure and evolution of the universe, life and entropy, nano-scale phenomena.					
<b>15. SUBJECT TERMS</b> EOARD, Modelling & Simulation, Quantum Chemistry, Nanotechnology					
<b>16. SECURITY CLASSIFICATION OF:</b>			<b>17. LIMITATION OF ABSTRACT</b> UL	<b>18. NUMBER OF PAGES</b>  60	<b>19a. NAME OF RESPONSIBLE PERSON</b> Ingrid Wysong
<b>a. REPORT</b> UNCLAS	<b>b. ABSTRACT</b> UNCLAS	<b>c. THIS PAGE</b> UNCLAS			<b>19b. TELEPHONE NUMBER (Include area code)</b> +44 (0)20 7514 4285

20030320 029

**Brijuni Conference VIII**

**Space, Time and Life**

**August 26 - 30, 2002**

**Brijuni, Croatia**

**Editor: S.D. Bosanac**

**NATO Advanced Research Workshop**

**Institut Rudjer Bošković  
Zagreb, 2002**

**ISBN 953-6690-23-3**

**We wish to thank the following for their contribution to the success of this conference:**

European Office of Aerospace Research and Development  
Air Force Office of Scientific Research  
United States Air Force Research Laboratory  
Ruđer Boskovic Institute  
Ministry of Science and Technology of Croatia  
Wolfram Research Inc.  
Systemcom d.o.o.

**Organizing committee:**

S. D. Bosanac - chairman  
H. Kroto - co-chairmen  
P. Toennies  
F. Scappini

**Local Organizing Committee:**

S.D. Bosanac  
J. Bosanac  
N. Došlić  
A. Mišević  
D. Plavšić

**Scientific Committee:**

S. D. Bosanac, Croatia  
T. Ebbesen, France  
D. Kleppner, USA  
W. Klemperer, USA  
H. Kroto, UK  
S. Leach, France  
G. Pichler, Croatia  
M. Quack, Switzerland  
F. Scappini, Italy  
P. Toennies, Germany  
N. Trinajstić, Croatia  
J. Weiner, France

## **Contents**

<b>1. Program.....</b>	<b>4</b>
<b>2. Abstracts.....</b>	<b>10</b>
<b>3. List of Participants.....</b>	<b>55</b>

**Monday, 26. August**

8.45-9.00

**S. D. Bosanac** - Introductory remarks

Chairmen: **J. Magueijo**

9.00-9.40

**K. Mainzer** - Matter, Life, and Information -  
Phase Transitions of the Universe

9.40-10.00

discussion

10.00-10.40

**I. Picek** - On the cosmic connection of  
fundamental particles and fields

10.40-11.00

discussion

11.00-11.20

coffee break

11.20-12.00

**M. Jura** - Are Earth-like planets going to be  
discovered soon?

12.00-12.20

discussion

**LUNCH AND BEACH DISCUSSIONS**

Chairmen: **I. Picek**

17.30-18.10

**L. Blitz** - The earliest stages of star formation

18.10-18.30

discussion

18.30-19.10

**J. Magueijo** - The life of the universe

19.10-19.30

discussion

Opening of poster session

Chairmen: **L. Klasinc**

**J. Belchior** : Applications of artificial neural  
networks for modelling controlled release  
systems

**T. Hrenar**: Molecular Dynamics Simulations of  
the infrared spectrum of benzaldehyde

**I. Ljubić**: CASSCF/CASPT2 study of the  
mechanism and kinetics of the gas-phase ozone  
additions to ethene, fluoroethene and  
chloroethene

**A. Khodin**: 3+1 is more than enough - particles  
in lower dimensions

21.00

**E. Prudnikov:** Self-organization from space to life

**V.Noethig-Laslo:** A spin labelling study of the hydrophobic barrier in liposomes

**D. Bonacci:** Laser control: analytical solution of the three level system dynamics

**V. Mikuta-Martinis:** Time operator in quantum mechanics

**D. Plavšić:** Modeling of boiling points of benzenoid hydrocarbons revisited

**D. Aumiler:** Determination of the atom number density in rubidium vapour

**V. Živčec:** High-pressure cesium discharge lamp

**G. Kovačević:** Proton transfer in acetylacetone: an ab-initio study

**S. Nikolić -** Nanotubes: Number of Kekulé Structures and Aromaticity

**L. Klasinc -** Measurement and modeling of radical scavenging by flavonoids

**R. M. Beuc -** Semiclassical approach to the description of collisionally induced optical transitions of diatoms

21.45

Evening discussion on Has the speed of modern communications changed our standards of scientific accuracy and ethical behaviour ?

Moderator: **G. Scoles**



**Tuesday, 27. August**

**Chairmen: W. Klemperer**

- |             |   |
|-------------|---|
| 9.00-9.40   | <b>M. Mayor</b> - Extrasolar planetary systems: From Individual Detections to Statistical Properties. |
| 9.40-10.00  | discussion  |
| 10.00-10.40 | <b>H. Svensmark</b> - Solar influence on Earth climate  |
| 10.40-11.00 | discussion  |
| 11.00-11.20 | coffee break  |
| 11.20-12.00 | <b>P. Thaddeus</b> - New interstellar and laboratory molecules  |
| 12.00-12.20 | discussion  |

**LUNCH AND BEACH DISCUSSIONS**

**Chairmen: M. Martinis**

- |             |   |
|-------------|---|
| 17.30-17.50 | <b>L. Pogosian</b> - Signatures of matter-antimatter asymmetry in the cosmic microwave background radiation?            |
| 17.50-18.10 | <b>P. P. Delsanto</b> - Bridging from micro to macro by means of mesoscopic simulations in physics and biology          |
| 18.10-18.30 | <b>R. Twarock</b> - New symmetries for fullerenes: new tools for the description of Carbon onions and Carbon nanotubes. |
| 18.30-18.50 | <b>M. Randić</b> - On characterization of 2-D proteomics maps   |

**Chairmen: S. D. Bosanac**

- |       |  |
|-------|--|
| 21.00 | <b>H. Kroto</b> - Science, a round peg in a Square World |
|-------|--|

Wednesday, 28. August

Chairmen: N. Trinajstić

9.00-9.40	<b>G. Arrhenius</b> - The emergence of life; geochemical, molecular biological and informational constraints on theory and experiment
9.40-10.00	discussion
10.00-10.40	<b>M. Eigen</b> - Natural selection as a phase transition in sequence space
10.40-11.00	discussion
11.00-11.20	coffee break
11.20-12.00	<b>J.-M. Lehn</b> - Steps towards complex matter: information, self-organization and adaptation in molecular and supramolecular systems
12.00-12.20	discussion
PICNIC	
Chairmen: <b>G. Arrhenius</b>	
21.00	<b>C. Cantor</b> - The future of life

**Thursday, 29. August**

**Chairmen: F. Scappini**

9.00-9.40	<b>M. Quack</b> - On the role of molecular space and time asymmetries in the origin of life.
9.40-10.00	discussion
10.00-10.40	<b>P. Walde</b> - Current ideas about the first cells and their precursor structures
10.40-11.00	discussion
11.00-11.20	coffee break
11.20-11.40	<b>S. Svetina</b> : Lipid vesicles and the origin of some cellular processes
11.40-12.00	<b>D. Juretić</b> - Life and entropy production
12.00-12.20	<b>E. Gallori</b> - Origin and persistence of genetic material in prebiotic habitats

**LUNCH AND BEACH DISCUSSIONS**

**Chairmen: P. Walde**

17.30-18.10	<b>S. Leach</b> - Panspermia: myth or reality?
18.10-18.30	discussion
18.30-19.10	<b>J. Durup</b> - The time variable in the neural encoding of consciousness states
19.10-19.30	discussion
20.30	banquet

**Friday, 30. August**

**Chairmen: G. Pichler**

9.00-9.40	<b>J. P. Toennies</b> - Diffraction of matter waves from nanostructures: new physics and novel applications
9.40-10.00	discussion
10.00-10.40	<b>T. Ebbesen</b> - Controlling optical diffraction and directionality from subwavelength apertures
10.40-11.00	discussion
11.00-11.20	coffee break
11.20-12.00	<b>W. Klemperer</b> - Intermolecular interactions
12.00-12.20	discussion
12.20	<b>M. Harwit</b> - Overview of the conference

## **2. Abstracts**

## **THE EMERGENCE OF LIFE; GEOCHEMICAL, MOLECULAR-BIOLOGICAL AND INFORMATIONAL CONSTRAINTS ON THEORY AND EXPERIMENT.**

Gustaf Arrhenius

Scripps Institution of Oceanography, University of California, San Diego, La Jolla, CA  
92093-0236

The questions concerning the origin and early evolution of life are mostly approached with focus on the complex problems of biomolecular autotynthesis. The specialists in this field are chemists and molecular biologists. They are seriously concerned with the staggering difficulties met in their own realm and tend to avoid dealing with the additional severe environmental limitations that are suggested by our general knowledge of planetary and spatial physics and chemistry. This purposeful neglect is often rationalized by the impression that their counterparts in the environmental sciences are groping in the dark and that, allowing enough time, shifting geochemical consensus will give back unlimited freedom to organic suppositions.

Planetary chemists on the other hand draw on their experience in studying present-day natural processes with reflection in the geological record. This makes them fit to attempt reconstruction of the environmental processes that may have controlled biogenesis. However, they often labor under the impression that any kind of organic molecule that can be conjured up in a harsh Archean environment signifies the emergence of life.

The gap between the two cultures has been narrowed, but not closed, by the extensive interdisciplinary activities in this field, fostered chiefly by NASA and other national and international agencies with an interest in exploring the origin of life and the possibility of its emergence also on other worlds. The present meeting gives an opportunity for another step toward defining the central problems and for reviewing relevant progress. This presentation will aim at the following central topics:

- " Environmental requirements for the generation of source molecules needed for metabolism and for generation and transduction of information
- " Factors in the primordial planetary and spatial environments that favor constructive processes vs. detrimental conditions
- " Physically and chemically realistic conjectures that can be made to overcome apparent hindrances
- " Autotrophic generation, encoding and propagation of molecules capable of information transduction
- " Modeling of primordial metabolism
- " Concentration and encapsulation of source molecules from highly dilute state; the improbability of a "primordial soup"
- " Earliest planetary evidence for life and its relation to the phylogenetic tree on Earth

## DETERMINATION OF THE ATOM NUMBER DENSITY IN RUBIDIUM VAPOUR

D. Aumiler, T. Ban, R. Beuc and G. Pichler

Institute of Physics, Bijenička 46, P.O. Box 304, HR-10001 Zagreb, Croatia

A new method for the atom number density determination in alkali vapour is proposed. The method is based on the comparison of numerical simulation of the alkali self-broadened resonance lines absorption profiles with measured ones. The numerical simulation calculates absorption profiles starting from the relevant set of alkali *ab initio* potential curves, using semiclassical stationary point approximation. As a result, the reduced absorption coefficient (absorption coefficient divided by the square of the atom number density) is obtained. By adjusting the temperature parameter of the model, it is shown that in the spectral region around resonance lines excellent agreement between simulated and experimental absorption profiles can be achieved. The least squares fitting of the simulated profiles to the measured profiles determines the vapour temperature with the uncertainty of less than  $\pm 1^\circ\text{C}$ . The measurements were made on rubidium vapour generated in a homemade crossed heat-pipe oven. In this way we were able to measure the Rb atom number densities with accuracy better than  $\pm 3\%$ .


The advantage of this method is that the measurement of the sole physical quantity – absorption coefficient is sufficient for the accurate determination of the atom number density. In particular, one does not need to know the temperature of the vapour in order to determine the atom number density. This is especially important in experiments in which exact vapour temperature measurements are not possible, as is the case when a heat-pipe oven is used. Similar application of the method could also be found in plasma diagnostics of pulsed high-pressure alkali light sources, which we are currently investigating in our laboratory.

## APPLICATIONS OF ARTIFICIAL NEURAL NETWORKS FOR MODELLING CONTROLLED RELEASE SYSTEMS.

Jadson C. Belchior\*

Departamento de Quimica - ICEX, Universidade Federal de Minas Gerais  
Pampulha, (31.270-901) Belo Horizonte, MC – Brazil

Efficient controlled release systems are fundamental for efficient therapeutical methodologies mainly to improve drug treatment through rate and time programmed drug delivery. However, the occurrence of multicomponents instead of single drug diffusion may complicate the experimental procedure to produce adequate drug delivery systems. Therefore, mathematical models of controlled delivery systems are important to simulate the device properties such as the geometries and physicochemical properties of the system.



This work presents an alternative methodology, based on artificial neural networks (see figure) to study drug delivery systems. A drug model system of hydrocortisone was selected to test the present approach but introducing modifications to increase complexity. The results of the neural network obtained after the learning stage can be considered quantitative to predict an ideal experimental condition. The analysis of the outcome results has shown that one can predict properties of an efficient experiment with a relative average error smaller than 2%.

Figure: Artificial neural network. Input layer (F - drug fraction released and t - time), an intermediary layer and an output layer (diffusion coefficient (D), saturation solubility (C<sub>i</sub>), initial drug load (A) and height of the device (h)).

\*Work supported by CAPES, FAPEMIG and CNPQ, Brazil.



## SEMICLASSICAL APPROACH TO THE DESCRIPTION OF COLLISIONALLY INDUCED OPTICAL TRANSITIONS OF DIATOMS

R. M. Beuc and M. Movre

Institute of Physics, Institute of Physics, Bijenička 46, P.O. Box 304  
HR-10001 Zagreb, Croatia

In a recent paper by Devdariani et al. [1], a formula was obtained that describes asymptotically forbidden quasimolecular optical transitions in the frame of the semiclassical approach. The main difficulty in the analytical description of asymptotically forbidden transitions is the consequence of the simple fact that the transition moments in this situation cannot be approximated by constant values as in the case of allowed transitions. However, potential energy curves and optical transition probability (or radiation width) usually can be reasonably approximated by exponential functions and a general formula which covers the central part of the spectral line, the extremum vicinity and the far wings, and also takes into account the fast exponential change in the state radiative width, can be obtained with the use of the Morse potential for the potential energy difference.

Our approach is based on a refinement of the theoretical model, based on the uniform Airy approximation as presented by Beuc and Horvatic [2]. The main point is careful treatment of the phases corresponding to the relevant Condon points and the resulting interference effects which in the same time take care of the variable transition moment in the vicinity of an extremum in the curve describing the difference between the electronic energies of the initial and the final states of the optical transition.

Bieniek et al. [3] have analyzed analogous processes of ionizing collisions in complex potentials within uniform JWKB stationary-phase techniques. Their numerical results are in excellent agreement with fully quantal, complex-potential computations. Our new results additionally support conclusions reached in [2] and [3] that the unified Franck-Condon theory is capable to describe asymptotically forbidden but collisionally induced transitions.

---

[1] A. Devdariani, E. Bichoutskaia, E. Tchesnokov, T. Bichoutskaia, D. S. F. Crothers, E. Leboucher-Dalimier, P. Sauvan and P. Angelo, *J. Phys. B: At. Mol. Opt. Phys.* **35** (2002) 2469-2475

[2] R. Beuc and V. Horvatic, *J. Phys. B: At. Mol. Opt. Phys.* **25** (1992) 1497-1510

[3] R. J. Bieniek, M. W. Müller and M. Movre, *J. Phys. B: At. Mol. Opt. Phys.* **23** (1990) 4521-4538

## **THE EARLIEST STAGES OF STAR FORMATION**

Leo Blitz

University of California at Berkeley, Radio Astronomy Laboratory, 635 Campbell  
3411, Berkeley, CA-94720-3411, USA

Serious searches for signals from technological civilizations will be greatly enhanced by a new generation of radio telescopes, the first of which is currently under construction by the Radio Astronomy Laboratory at Berkeley. The probability that the searches will be successful depends, among many other things, on how commonly the conditions conducive to the development of life are found in the Milky Way. The birth of stars and planetary systems takes place in giant clouds of molecular gas and dust which are remarkably similar to one another throughout the Milky Way and in nearby galaxies. The sun-like stars that form within them inevitably have protoplanetary disks that have a large range in properties and are affected by environmental differences. If biological evolution requires catastrophic events such as bombardment by cosmic debris, the evidence suggests that although planetary systems vary greatly, such debris is common. Its distribution, and thus the likelihood of colliding with an earth-like planet, depends on the details of planet formation. Although the formation of earth-like systems is probably not as common as previously thought, recent evidence suggests that the differences between systems are less important than their similarities.

## **LASER CONTROL: ANALITICAL SOLUTION OF THE THREE LEVEL SYSTEM DYNAMICS**

D. Bonacci, S.D. Bosanac, N. Došlić  
'Ruđer Bošković' Institute, Bijenička 54, 10000 Zagreb

Many systems of current interest in physical chemistry are too complex for the full analytical treatment. For example, laser control parameters for manipulation of molecular state population are generally optimized numerically. Although the advancement of computer technology eliminated the time consuming note from this approach, it still lacks the depth of theoretical insight that even approximate analytical calculation would yield.

We obtained analytical solution of the specific three-level system in strong coherent electric field, extending in this way the classical Rabi theory developed half a century ago for two-level systems. Analytical solution enabled us to explore some details of the system's dynamics, and to determine the relative importance of various physical parameters of the system.

## **THE FUTURE OF LIFE**

Charles R. Cantor, PhD  
SEQUENOM, Inc., San Diego, CA USA

All living organisms have properties that are a convolution of their genetic programming and environmental exposure. Until recently humankind has had only a feeble ability to modify the genetic programming of organisms, but in recent years this has changed dramatically. We now can pretty much do anything we want. However, our current ability to predict the outcome of genetic manipulations is still extremely limited. In parallel, humans have also gained the ability to manipulate the environment. While we are far away from total control, globally, we can do pretty well, locally.

The result of these advances in technology is that future evolution of all life on our planet will be pretty much controlled by humans. We will see the emergence of newly designed species with enhanced properties to communicate, regenerate, produce energy efficiently, generate fuels, synthesize medicines, and manufacture healthier (for humans) food. We will also witness the directed evolution of humans with enhanced properties. Clearly this will not occur without some controversy.

It is probably a natural outcome of evolution that a species attains the ability to control and accelerate its own evolution. However, this great power to improve also comes with an equal power to do harm, and no single technology has greater power to threaten our stability as a society than biotechnology, if used malevolently.

## **BRIDGING FROM MICRO TO MACRO BY MEANS OF MESOSCOPIC SIMULATIONS IN PHYSICS AND BIOLOGY**

Pier Paolo Delsanto

Dipartimento di Fisica, Politecnico di Torino, C. Duca Abruzzi 24, Torino 10129  
ITALY

An impressive range of results has been obtained in the field of Molecular Dynamics in the last decade for systems up to the size of about 100 nm. Due to the enormous potential for technological applications at the macroscopic level it is necessary to go well beyond such a limit. However, the computational costs of increasing the size of the system of even one or two orders of magnitude are absolutely prohibitive. A suitable alternative may be to split up the problem according to the different length scales involved, i.e. Multiscale Modelling (MM). Several approaches have been proposed for MM: in general they start from atomistic simulations, replacing, however, the interatomic interactions with e.g. interactions among dislocations or, for larger systems, among dislocation densities or grains. However, since our major concern lies in the macroscopic domain (e.g. NDE or materials characterization), we propose to reverse this "bottom-up" strategy.

The approach we propose starts with a very fine discretization of the macroscopic specimen under study, which brings us to the mesoscopic level. For this bridging from a microscopic to a macroscopic scale by means of mesoscopic simulations, we plan to use the local Interaction Simulation Approach (LISA), whose "local" nature makes it ideally suitable for the task in hand, particularly in the case of strong local heterogeneities. Due to the complexity and vastness of scope of our goal, rather than searching for a general recipe, we propose to investigate several "case studies", which should demonstrate the feasibility of the approach and its validity. A similar approach can be used in biology and medicine to bridge between cell dynamics and macroscopic problems, such as the simulation of growth and therapy of tumors.

## PROTON TRANSFER IN ACETYLACETONE: AN AB-INITIO STUDY

G. Kovačević, I Matanović and N. Došlić  
R. Bošković Institute, Bijenička 54, Zagreb Croatia

Reaction minima and transition states, concerning proton transfer in acetylacetone were determined using Moller-Plesset ab initio and B1LYP density functional theory calculations. It is found that the minima and the transition state for proton transfer have different methyl groups orientation. Intrinsic reaction path calculation has shown that the rotation of the methyl groups is not concerted with the proton transfer reaction, rather they are consecutive processes. The thermodynamical stability of the keto and enol forms of acetylacetone is explored by using G2 type calculations and it is shown to be in good agreement with experimental results.

## THE TIME VARIABLE IN THE NEURAL ENCODING OF CONSCIOUSNESS STATES

Jean Durup

Laboratoire de Physique Quantique (Université Paul Sabatier et CNRS UMR 5626),  
IRSAMC, 118 route de Narbonne, 31062 Toulouse, France

That *time* plays an important role in the so-called NCC (Neural Correlate of Consciousness) has been suspected for long time, both because *synchrony* appeared to be critical for the binding of the various features building a conscious representation, and from the experimental observation of stronger *oscillations* of the electric field associated with the performance of a cognitive task when performed in a conscious way.

Nevertheless, although probably 'in the air' for more than a decade, the simple theory claiming that consciousness is characterized by the encoding of its content into a *purely temporal pattern*, irrespectively of the identity of the neurons involved, has been either ignored or rejected, although having been the subject of 3 recent works (Helekar 1999, Durup and Sanejouand 2000, John 2001).

We shall show that the combination of largely accepted experimental data with insights from evolution theory allows us to give a *demonstration* of the above-stated theory, on the basis of the immense ratio between the capacities of non-conscious vs conscious processing.

Important differences between the models which have been developed within this theory will be discussed: (i) is the code universal or individual ? (ii) are the relevant neural computations performed with bits or with qubits ?

Finally, and in an unrelated way, we will show that a well-known aspect of consciousness, namely its limited content, allows us to give an elegant answer to the old philosophical problem: free will or determinism ?

## **CONTROLLING OPTICAL DIFFRACTION AND DIRECTIONALITY FROM A SUBWAVELENGTH APERTURE**

Thomas W. Ebbesen

ISIS, Louis Pasteur University, 4 rue B. Pascal, 67000 Strasbourg, France

Materials structured on the nanometer scale can lead to improved and sometimes surprising properties. This will be illustrated in some detail by our own work on the extraordinary optical properties acquired by metal films when their surfaces are corrugated and perforated with one or more sub-wavelength holes ( $\sim 150$  nm). Such structures can transmit the light with an efficiency hundred times larger than what theory predicts for single holes. The efficiency can even be much larger than the fractional area occupied by the hole, which means that even the light falling beside the hole emerges on the other side of the sample. This extraordinary transmission is due to the coupling of the incident light with the surface plasmons of the film. The transmission spectrum contains peaks attributed to surface-plasmon modes that depend on both the symmetry and the 2D lattice parameter of the surface corrugation. Novel results show that another fundamental problem of sub-wavelength apertures, namely optical diffraction, can also be modified using surface plasmons. The optical divergence and directionality of light transmitted by a single subwavelength aperture can be controlled. These results have broad fundamental and practical implications and show that, with modern fabrication techniques, surface plasmons can be engineered to yield unique optical properties.

Ref.: Ebbesen et al Nature 391, 667 (1998); Ghaemi et al, Phys. Rev. B 58, 6779 (1998); Grupp et al Adv. Mater. 11, 860 (1999); Kim et al, Optics Letters 24, 256 (1999); Grupp et al, Appl. Phys. Lett., 77, 1569 (2000); Martin-Moreno et al, Phys. Rev. Lett. 86, 1114 (2001); Krishnan et al. Optics Comm. 200, 1 (2001); Lezec et al Science, in press (2002).



## BIOLOGICAL INFORMATION AND PHASE TRANSITIONS

M. Eigen

MPI fuer Biophysikalische Chemie, Goettingen

Information has two aspects: a quantity to be called "extent" and a quality which may be termed "content" since it deals with meaning. The latter originates via selective self-organization which can be described also in quantitative physical terms. A prerequisite is the reproducibility of the informational substrate forming the basis of selection. This paper focuses on selection being the analogue of a physical phase transition. In the first part the criteria for phase transitions are formulated. The second part introduces the concept of information space and describes information as selected points or regions in this space. In the third part selection is analysed in terms of the criteria for phase transitions, and finally the concept is confronted with experimental data. The conclusion is reached that information content is generated via selection which can be described as a phase transition in information space.

## **ORIGIN AND PERSISTENCE OF GENETIC MATERIAL IN PREBIOTIC HABITATS**

Enzo Gallori

Department of Animal Biology and Genetics, University of Florence, Via Romana  
17/19, 50125 Florence, Italy

Molecules which store genetic information (RNA, DNA) are central to all life on earth. The formation of these complex molecules required specific conditions, including the synthesis of precursors (nucleotides), the joining of these monomers into larger molecules (polynucleotides), their protection from degradation, and the expression of the biological potential of the informative molecule, which is its capacity to multiply and evolve. Determining how these steps occurred and how the earliest genetic molecules originated on Earth is a problem that is far from being resolved. Classical research in this field has focused on process in aqueous solution. However, it is difficult to believe that complex molecules can be obtained by random collisions in a fully aqueous environment since under these conditions, hydrolysis and not polymerization is favoured.

We believe that it is crucial to consider surface organic chemistry.

As early as 1951, J.D. Bernal suggested, on the basis of thermodynamic and kinetic considerations, that mineral surfaces (specifically clay minerals) could have played an important role in the prebiotic formation of the biomolecules basic to life.

In recent years, numerous observations in different fields, from astrophysical chemistry to molecular biology, have reinforced the hypothesis of the involvement of mineral surfaces in the prebiotic origin of life.

Brijuni Conference VIII, Brijuni, Croatia, 26.8.-31.8.2002.

## **OVERVIEW OF THE CONFERENCE**

Martin Harwit

Cornell University, Department of Astronomy, Ithaca, NY 14853-6801, USA

This conference covers the entire range of issues --- cosmological, astronomical, geophysical, chemical, biological, and neurological --- dealing with the questions "How did we get here? How is it that we are even able to ask this question?" This final talk will attempt to summarize what the previous speakers have taught us and to assemble their thoughts into a coherent thread.

## **MOLECULAR DYNAMICS SIMULATIONS OF THE INFRARED SPECTRUM OF BENZALDEHYDE**

T. Hrenar(1), R. Mitrić(2) and Z. Meić(1)

(1) Faculty of Science, University of Zagreb, Strossmayerov trg 14, HR-10000  
Zagreb, Croatia

(2) Humboldt Universität zu Berlin, Institut für Chemie, Brook-Taylor-Str. 2 12489  
Berlin

A calculation of vibrational spectra with semiempirical or ab initio force fields gives generally too high values for vibrational frequencies. However, the use of the density functional methods (DFT) reduces the need for correction of frequencies or the corresponding force constants. On the other hand, calculations of vibrational spectra by molecular dynamics methods do not demand such empirical corrections.

With the calculated ab initio force fields we have carried out Newtonian dynamics. From the time dependency of the dipole moment one can deduce information on infrared (IR) spectra of molecules. The described method is applied to benzaldehyde, which is a proper molecule due to its size and availability of experimental data. The obtained results will be presented and discussed.

## **ARE EARTH-LIKE PLANETS GOING TO BE DISCOVERED SOON?**

Michael Jura

Department of Physics and Astronomy, University of California, Box 951547, Los Angeles, CA 90095-1547, USA

Jovian-mass planets and "pebbles" as large as 5 cm in diameter have been inferred to orbit nearby stars. However, to date, there has been no direct evidence for the existence of terrestrial-mass planets, although we strongly suspect that such planets exist.

In this talk, the evidence for the existence of extra solar-system asteroids will be reviewed, and strategies for the discovery of terrestrial mass planets will be discussed. Methods to search for life on these hypothetical planets are part of this program.

## **LIFE AND ENTROPY PRODUCTION**

Davor Juretić

Department of Physics, University of Split, Nikole Tesle 12, Split, Croatia

Nobody disputes that life creates entropy, but whether living systems augment or diminish the rate of entropy production is still an unresolved question. Do living entities make the least concession possible to entropy by seeking the steady state of minimal entropy production, or do they maximize the rates of dissipation?

We shall use several simple models for cyclic biochemical reactions to calculate entropy production and efficiency of free-energy conversion in a steady state. Our goal will be to explore whether actual operation of these cycles can be better modeled by requiring minimal or maximal entropy production. An optimization principle coming from physics would surely help in predicting behavior of complex metabolic networks revealed after the completion of different genome projects.

Main attention will be devoted to bioenergetics of bacterial and plant photosynthesis, and to ATP driven biochemical cycles.

### 3+1 IS MORE THAN ENOUGH – PARTICLES IN LOWER DIMENSIONS

A.Khodin

National Academy of Sciences of Belarus, Minsk

The topological approach is developed where an elementary particle is considered as corresponding to a localised space-time configuration [1]. The key of the approach is the standard relativistic limit for space-time curvature. In general case, the space-time configurations, for which this limit is maintained, are shown to be three-, two-, and one-dimensional as space objects. The real space-time is proposed to consider as evolution of a mixture of one-, two-, and three-dimensional micro-objects, which interacts and compose macroobjects corresponding to its degrees of freedom in space co-ordinates.

In the frames of the approach presented, a photon is described as 2+1 configuration where abovementioned condition is valid in its centre point. The configuration possesses of Doppler effect behaviour in the case it propagates with light velocity only. Topological charge of the configuration is zero. The photon spin is represented as configuration' rotation around the propagation direction; the only permissible rotation period corresponding to the photon wavelength. It is shown that ordered one-dimensional superposition of single-photon configurations composes a restricted stable coherent harmonic-type packet which wavelength equals to the constitutive photon configurations' wavelength. Two- and three-dimensional coherent superpositions of single-photon configurations are shown to form some types of photons ordering symmetry inside the coherent packet.

1. A.Khodin. EPS-11: Trends in Physics. 11th Gen. Conf. of Europ. Phys. Soc. London, UK, 6-10 Sept.1999.

## INTERMOLECULAR FORCES

William Klemperer

Harvard University, Department of Chemistry, 12 Oxford Street, Cambridge, Mass.  
02138, USA

The characterization, at high spectral resolution, of molecular complexes has led to detailed knowledge of intermolecular forces. The structure and dynamics of stable and metastable states will be discussed. Of particular interest will be hydrogen bonded species as well as the rich variety of isomeric forms presently observed. We examine the question of whether and when weak interactions can modify dynamical pathways. Finally we discuss the structure of complexes of molecules with hydrogen and helium.



## **PANSPERMIA: MYTH OR REALITY?**

Sydney Leach,

Observatoire de Paris-Meudon, 92195 - Meudon, France

Panspermia, the hypothesis that life reached Earth from outer space, was developed by Svente Arrhenius between 1903 and 1906. This is the "strong" version of panspermia, and it sidesteps the question of how did life originate. There were several precursors who had suggested that life existed elsewhere in the Universe (Anaxagoras, Giordano Bruno, Fontanelle). There were also precursors such as Berzelius who, in 1834, suggested meteorites as carriers of life. But a scientific approach to these questions had to await the demise of the long-held concept of spontaneous generation and the development of Darwin's theory of evolution. As understanding of astronomical and astrophysical observations progressed, the concept of life elsewhere than on Earth had fluctuating fortunes, as did estimations of the probability of survival of space transit and atmospheric entry. Thus a more chemical/biochemical approach to the origin of life on Earth developed, one variant of which, involving the infall of prebiotic chemicals, constitutes a "weak" panspermia hypothesis. The scientific and, in part, philosophical background to these developments will be discussed and will include remarks on directed panspermia.

## STEPS TOWARDS COMPLEX MATTER: INFORMATION, SELF-ORGANIZATION AND ADAPTATION IN MOLECULAR AND SUPRAMOLECULAR SYSTEMS

Jean-Marie LEHN

ISIS, Université Louis Pasteur, Strasbourg and Collège de France, Paris

One of the major lines of development of chemical science in the coming century resides in the ever clearer perception, deeper analysis and more deliberate application of information features in the elaboration and transformation of matter, thus tracing the path from merely condensed matter to more and more highly organized matter towards systems of increasing complexity.

A most basic and far-reaching contribution of supramolecular chemistry is the implementation in chemical science of the concept of molecular information and its corollaries, instructions and programmed chemical systems, with the aim of gaining progressive control over the organization of matter, over its spatial (structural) and temporal (dynamical) features.

Supramolecular chemistry is actively exploring systems undergoing *self-organization*, i.e. systems capable of spontaneously generating well-defined functional supramolecular architectures by self-assembly from their components. Self-organization processes may be directed via the *molecular information* stored in the covalent framework of the components and read out at the supramolecular level through specific interactional algorithms, thus behaving as *programmed chemical systems*.

The control provided by self-organization processes allows the development of *functional supramolecular devices*, defined as structurally organized and functionally integrated systems built from suitably designed molecular components performing a given function (e.g. photoactive, electroactive, ionoactive, etc.) and containing the features required for assembly into an organized supramolecular architecture. Self-organization processes also give access to advanced *functional supramolecular materials*, such as supramolecular polymers and liquid crystals.

The design of molecular information controlled, "*programmed*" self-organising systems provides an original approach to *nanoscience* and *nanotechnology*. In particular, the spontaneous but controlled generation of well-defined, functional supramolecular architectures of nanometric size through self-organization represents a means of performing programmed *engineering* and *processing* of *nanomaterials*.

Technologies resorting to self-organization processes are in principle able to provide a powerful alternative to nanofabrication and nanomanipulation procedures by making use of the spontaneous but controlled generation of the desired superstructures and devices from suitably instructed and functional building blocks.

Supramolecular chemistry is also intrinsically a *dynamic chemistry* in view of the lability of the interactions connecting the molecular components of a supramolecular entity. Furthermore, because of the ability of supramolecular species to exchange their constituents, they have also *combinatorial* capacity.

Consequently, supramolecular devices and materials are by nature *dynamic species*, whose constituents are linked through reversible connections and undergo spontaneous and continuous assembly/deassembly processes in a given set of conditions. Because of their intrinsic ability to exchange their constituents, they have also combinatorial character so that they may be considered as *dynamic combinatorial devices and materials*. Being instructed, dynamic and combinatorial, they may in principle select their constituents in response to external stimuli or environmental factors and therefore behave as *adaptive systems*.

The same considerations apply to molecular entities constructed from components linked through reversible covalent connections. Together, such molecular and supramolecular systems embody a *dynamic combinatorial chemistry*.

The merging of the features of supramolecular systems : - information and programmability, - dynamics and reversibility, - constitution and structural diversity, points towards the emergence of *adaptive chemistry*. A further development will concern the inclusion of the arrow of time, i.e. of non-equilibrium, irreversible processes and the exploration of the frontiers of chemical evolution towards the establishment of *evolutionary chemistry*, where the features acquired by adaptation are conserved and transmitted. In combination with the corresponding fields of physics and biology, chemistry thus plays a major role in the progressive elaboration of a science of informed, organized, evolutionary matter, a *science of complex matter*.

#### *General references*

- J.-M. LEHN, *Supramolecular Chemistry: Concepts and Perspectives*, VCH, Weinheim, 1995.
- J.-M. LEHN in *Supramolecular Chemistry : Where It is and Where It is Going* (R. Ungaro, E. Dalcanale, eds.), Kluwer Dordrecht, 1999, pp.287-304.
- J.-M. LEHN, *Chem. Eur. J.*, 1999, 5, 2455.
- J.-M. LEHN, *Chem. Eur. J.*, 2000, 6, 2097.
- J.-M. LEHN, *Proc. Natl. Acad. Sci. USA*, 2002, in press.

**"CASSCF/CASPT2 STUDY OF THE MECHANISM AND KINETICS OF THE  
GAS-PHASE OZONE ADDITIONS TO ETHENE, FLUOROETHENE AND  
CHLOROETHENE".**

Ivan Ljubić

R. Bošković Institute, Bijenička 54, Zagreb, Croatia

"Ab initio multiconfigurational CASSCF and CASPT2 methods were employed in studying the reaction mechanisms and kinetics of the gas-phase ozone additions to ethene, fluoroethene and chloroethene. Reactants, transition state structures and products were optimized and harmonic vibrational frequencies calculated at the CASSCF/cc-pVTZ level. For kinetic calculations the electron energies of all the stationary points were further refined by utilizing the CASPT2 method with the optimized CASSCF/cc-pVTZ wavefunctions taken as the zeroth-order. The rate constants and Arrhenius kinetic parameters were finally calculated in terms of the conventional transition state theory. The favourable conformations of the ozone approach to the two asymmetrically substituted haloalkenes are at first governed by the electrostatic repulsion in the transition state structures and later by the gradually predominating anomeric effect. The bond formation in the primary haloozonides was analyzed by monitoring the changes in the occupation numbers of the active orbitals in course of the optimizations. For all the reactions thus studied a close agreement is found with the experimental kinetics, which makes a future use of the same approach very promising."

## **THE LIFE OF THE UNIVERSE**

Joao Magueijo

Imperial College of Science, Technology and Medicine, Department of Physics,  
Exhibition Road, London SW7 2AZ, UK

Many results on the probability of life in the Universe assume that physics is close to a final answer, in particular that the theory of relativity is, at least, very approximately correct. But new astronomical observations and theoretical work suggest otherwise. Perhaps the Universe changes its laws as it expands, the constants of nature are wildly different near black holes and other high concentrations of matter, and ultimately the "nothing can travel faster than light" dogma is not entirely correct. I examine these possibilities with reference to what they may imply to life in the Universe, and our prospects of ever establishing communication with alien civilizations.

**MATTER, LIFE, AND INFORMATION  
PHASE TRANSITIONS OF THE UNIVERSE**

Klaus Mainzer

Universität Augsburg, Universität Str. 10, D-86159 Augsburg, Germany

In classical physics, space, time, and matter were considered as basic concepts of a steady-state universe. In the framework of modern science, the universe is no longer a dead mechanical clockwork: It seems to evolve as a self-organizing complex system according to phase transitions with the emergence of new materials, life, consciousness, and intelligence. Life is no random event at the edge of the universe. Are we alone? What are the common structures and laws of this development? In the age of computers, computer simulations and computer experiments deliver first insights into the information processing of complex systems. Information itself seems to be a fundamental ingredient in the evolution of the universe. These concepts of modern science lead back to ancient ideas in the philosophy of nature.

*Literature:* K. Mainzer, *Symmetries of Nature*, Berlin: De Gruyter 1996; *Thinking in Complexity: The Complex Dynamics of Matter, Mind, and Mankind*, New York: Springer 1997 (3<sup>rd</sup> ed.); *The Little Book of Time*, New York: Copernicus Books 2002.

## **EXTRASOLAR PLANETARY SYSTEMS: FROM INDIVIDUAL DETECTIONS TO STATISTICAL PROPERTIES**

Michel Mayor

Observatoire de Geneve, Switzerland

The discovery of planets around other stars was (up to now) only possible due to the development of high-precision radial-velocity techniques. These techniques have permitted astronomers to look for the tiny periodic motion of a star as it moves around the center-of-mass of the star-planet system. Radial velocity surveys have revealed up to now about 100 extra-solar planets and 9 multi-planetary systems. The discovered planets present a wide variety of orbital element and masses, which are raising many problems and questions regarding the processes involved in their formation. But the analysis of the distributions of orbital elements, like the period and eccentricity distributions is already giving some constraints on the formation of planetary systems. Furthermore, the study of planet host stars have revealed the impressive role of the stellar metallicity on the giant planet formation. The chemical composition of the molecular cloud is probably one of the key parameters to form giant planets. The study of extra-solar planetary systems is just giving its first steps. After only 7 years, we can say that at least 5% of the solar type stars have giant planetary companions with masses as low as the mass of Saturn and orbital periods less than about 5 years. But the understanding of how giant planets are formed is still shaded in many points and we are still lacking of any detections of analogs of terrestrial planets. To help solve some of these problems, several projects are currently in the pipeline. Future space missions, like the photometric missions COROT, Eddington or Kepler (to search for planetary transits) or the astrometric satellites GAIA and SIM will definitively permit to unveil hundreds (or even thousands) of "new" planets. But ground based astronomy will also give enormous steps in the next few years.

## TIME OPERATOR IN QUANTUM MECHANICS

V. Mikuta-Martiniš (1) and M. Martiniš (2)

(1) "Rudjer Boskovic" Institute, Zagreb, Croatia,

(2) "Rudjer Boskovic" Institute, Zagreb, Croatia,

Controversy about the existence of a self-adjoint time operator,  $T$ , conjugate to a given Hamiltonian is reexamined. Specifically, we attempt to obtain an explicit solution to the operator equation  $-i [T(p,q), H(p,q)] = 1$  for a conformally invariant Hamiltonian. References: 1. V. de Alfaro, S. Fubini, G. Furlan: *Nuovo Cimento* 34A, 569 (1976). 2. C.M. Bender, G.V. Dunne: *Phys.Rev. D*40, 2739 (1989). 3. H.R. Lewis, W.E. Lawrence, J.D. Harris: *Phys. Rev. Lett.* 77, 5157 (1996).



## NANOTUBES: NUMBER OF KEKULÉ STRUCTURES AND AROMATICITY

István LUKOVITS,<sup>a</sup> Sonja NIKOLIĆ,<sup>b</sup> and Nenad TRINAJSTIĆ<sup>b</sup>

<sup>a</sup>Chemical Research Center, Hungarian Academy of Sciences, H-1525 Budapest,  
P.O.Box 17, Hungary

<sup>b</sup>Rugjer Bošković Institute, HR-10002 Zagreb, P.O.Box 180, Croatia

Carbon nanotubes (CNTs) are composed of cylindrical graphite sheets consisting of  $sp^2$  carbons. Due to their structure CNTs are considered to be aromatic systems. In this work the number of Kekulé structures (K) in CNTs was estimated by using the transfer matrix technique. All Kekulé structures of the cyclic variants of naphthalene and benzo[c]phenanthrene have been generated and the basic patterns have been obtained. From this information the elements of the transfer matrix were derived. The results obtained indicate that K (and the resonance energy) is greater if tubulenes are extended in vertical than in horizontal direction. Tubulenes are therefore more stable than cyclic strips.

## A SPIN LABELLING STUDY OF THE HYDROPHOBIC BARRIER IN LIPOSOMES

V.Noethig-Laslo<sup>a</sup>, P. Cevc<sup>b</sup>, D. Arcon<sup>b</sup> and M. Šentjurc<sup>b</sup>

<sup>a</sup> "R. Boskovic" Institute, 10002 Zagreb, Croatia,

<sup>b</sup> "J. Stefan" Institue, 10001 Ljubljana, Slovenia

A fundamental property of biological membranes is to act as a barrier to permeation of polar molecules. This barrier effect is largely due to a hydrophobicity of a phospholipid bilayer, determined by the extent of water penetration into the membrane.

In order to study the depth of water peneration into liposomes, we used the electron spin echo modulation, a pulsed electron spin resonance method [1], that takes advantage of the solvent effect on the hyperfine interaction of the nitroxide spin labels. A fatty acid with the paramagnetic nitroxide group of the spin label covalently attached at various positions along the fatty acid chain was introduced into the multilamellar liposomes. The hydrophobic barrier for water molecules, as well as, for small molecules such as glycine, was studied as a function of the liposome composition and size.

[1] A. Schweiger (1991) *Angew. Chem. Int. Ed. Engl.* **30**, 265-292.

## **ON THE COSMIC CONNECTION OF FUNDAMENTAL PARTICLES AND FIELDS**

Ivica Picek

Odsjek teorijske fizike, Bijenička c. 32, POB 162, 10 000 Zagreb, Croatia

We observe that, on the scene of Space and Time, the Matter arranges itself in ways that provide a chain of coincidences enabling creation of Life and intelligent creatures. While the broadest principles of Special Relativity and Quantum Mechanics provide merely a language, a further insight into the selection of particles and forces existing in the universe comes from the local Gauge Symmetries. The other cherished symmetries of the particle physics are awaited to be violated towards the more fundamental scale, probed in the early universe.

## MODELING OF BOILING POINTS OF BENZENOID HYDROCARBONS REVISITED

Dejan Plavšić,<sup>a</sup> Milan Šoškić,<sup>b</sup> Nella Lerš,<sup>a</sup> and Leo Frkanec<sup>a</sup>

<sup>a</sup>The Ruđer Bošković Institute, P.O.Box 180, HR-10002 Zagreb, The Republic of Croatia

<sup>b</sup>Faculty of Agriculture, University of Zagreb, HR-10000 Zagreb, The Republic of Croatia

Polycyclic aromatic hydrocarbons and their subset benzenoid hydrocarbons are a class of mutagenic and/or carcinogenic environmental pollutants being permanently formed by incomplete combustions. As benzenoids are ubiquitous pollutants new information about their properties and features as well as their biological activity are desirable and important.

The normal boiling point *per se* is an important feature of a compound, moreover it is one of the indispensable physicochemical properties for a reasonable risk assessment of a chemical. Hence, several QSPR (quantitative structure property relationship) studies of the normal boiling points of smaller benzenoid hydrocarbons have recently been carried out. Many of the normal boiling points used in the training set in all these studies differ considerably from the corresponding data in the sources like Beilstein, Merck Index, and others. For example, the normal boiling point of coronene in these studies is 590 °C, but according to the aforementioned reliable sources it is 525 °C!

We have carried out a novel QSPR study of the normal boiling points of benzenoid hydrocarbons using data from Beilstein and different topological indices as predictor variables. The developed models are mutually compared and some of them are of high quality.

Brijuni Conference VIII, Brijuni, Croatia, 26.8.-31.8.2002.

**“SIGNATURES OF MATTER-ANTIMATTER ASYMMETRY IN THE  
COSMIC MICROWAVE BACKGROUND RADIATION?”**

Levon Pogosian

Blackett Laboratory, Imperial College, Prince Consort Road, London, SW7 2BW,  
U.K.

Helical primordial magnetic fields could be expected from cosmic events such as electroweak baryogenesis. Such fields can lead to helical velocity flows and could manifest themselves as non-vanishing correlations of the temperature and B-type polarization of the CMBR. Primordial helical magnetic field can also be detected using Faraday rotation measure maps of the CMBR.

## SELF-ORGANIZATION FROM SPACE TO LIFE

Evgeniy D. Prudnikov

St. Petersburg State University, University emb. 7/9, St. Petersburg, 199034, Russia

Self-organization laws determine all natural processes from Space to Life. It is known the role of the fluctuations for the transition from dissipative structures to order (I. Prigogine). In the paper (Vestn. Leningrad Univ. 1991, ser. Physics, n. 25, P. 38-43), Prudnikov and Shapkina have proposed the description of the processes of self-organization on the fluctuation level. According to this criterion the level of self-organization of natural process may be presented by the following manner:

$$L_s = [l_0^2 + (k_d + 2RAI_0)S_i n + A^2 S_i^2 n^2 + k_k S_i^3 n^3 + k_n S_i^4 n^4]^{1/2} \quad (1)$$

Where  $L_s$  is the absolute level of self-organization of the given natural process, or the random fluctuations in this process;  $l_0$  is the zero level of self-organization, or the fluctuations for other processes of this system;  $S_i$  is the sensitivity of the process to the different influences, or the instability of the process, which is determined by the theory of the examined natural process and includes the flow of the energy in the system;  $n$  is the scale of the process, which is characterized by the quantity of the substance in this process or the substance concentration;  $k_d$ ,  $A$ ,  $k_k$  and  $k_n$  are the coefficients of the individual levels of the self-organization;  $R$  is the coefficient of the co-correlation. According to the equation (1), the value of  $S_i^{-1}$  determines the stability of the system in the conditions of the different influences. The equation (1) demands to observe the five levels of the fundamental relations and the states for the natural processes. These levels are connected with the stability of the process and its energy in the power 0,  $1/2$ , 1,  $3/2$ , 2. The first three terms on the right-hand side of above expression characterize the linear range of the dependence of the parameters of the process from the energy and scale of the process and system, whereas the following two terms determine the non-linearity with the change of the scale of the system.

One can distinguish the five basic levels of the conditions of the substance: submicro up to elementary particles; micro up to atoms; the linear conditions of the environment up to planets; the nonlinear world from stars up to black holes and hyper nonlinear world up to great explosion. The each natural process has also the five levels of the passing from zero up to stochastic, linear, nonlinear and catastrophic level. In reason of this the development of any process from space to life passes of the stochastic, linear and nonlinear stages in the dependence of the energy and the scale of the system, and is finished the transition in new condition of the system or the catastrophe. The evolution is the continuous chain of the development process from space up to intelligent life. The time ( $t$ ) is the function of the function of the speed of the process passing:  $t=F(S_i n)=F(L_s)$  (2). The quantitative characteristic of the self-organization processes and their comparison may be realized most effective by use of the relative value of the level of the self-organization ( $L_{sr}$ ):

$$L_{sr} = [l_0^2 S_i^{-2} n^{-2} + (k_d + 2RA l_0) S_i^{-1} n^{-1} + A + k_n S_i n + k_k S_i^2 n^2]^{1/2} \quad (3)$$

The expression (3) allows comparing the any natural processes according to the self-organization level independently the type and the theoretical basis of the processes. Here the time is connected with coefficient  $A$ , and the time is one order of the magnitude for any natural process. The theory allows tracing the evolution from space to life, from the simple systems up to complex systems, from to biological matter up to intelligent life. According to these positions the theory have not only the scientific importance, but the theory may play the essential philosophical role.

## ON THE ROLE OF MOLECULAR SPACE AND TIME ASYMMETRIES IN THE ORIGIN OF LIFE

Martin Quack

Physical Chemistry, ETH Zürich, CH-8093 Zürich, Switzerland

In the lecture we shall explore the role of the fundamental symmetries of physics in the molecular origin of life – in particular the space, time and matter – antimatter symmetries expressed in C, P, T symmetries and their combinations (CP, CPT). In particular, molecular and biomolecular chirality are related to these symmetries and may be considered to be molecular “fossils” of evolution. The two key concepts here are the tunnel effect (introduced by F. Hund for the stereomutation of chiral molecules in 1927 [1]) and the influence of the electroweak parity violation suggested already by Yamagata in 1966 to be of potential relevance for biomolecular chirality [2]. In the lecture we will mainly report about the recent efforts of the Zürich group related to the theoretical discovery (in 1996 [3]) that parity violating electroweak potentials are orders of magnitude larger than previously calculated. We will provide the general theoretical background [4-6] as well as recent applications [7-10]. We shall mention current spectroscopic efforts on molecular parity violation [11-13], as well as possible conclusions to be drawn for the biomolecular evolution of life [14-16].

- [1] F. Hund, *Z. Phys.*, **1927**, *43*, p. 788-844, 805-826.
- [2] Y. Yamagata, *J. Theor. Biol.*, **1966**, p. 495-498.
- [3] A. Bakasov, T. K. Ha, M. Quack, "Ab initio calculation of molecular energies including parity violating interactions", p. 287-296, in "Chemical Evolution, Physics of the Origin and Evolution of Life" (Eds.: J. Chela-Flores, F. Raulin), Kluwer Academic Publishers, Dordrecht, **1996**
- [4] A. Bakasov, T. K. Ha, M. Quack, *J. Chem. Phys.*, **1998**, *109*, p. 7263-7285.
- [5] M. Quack, *Nova Acta Leopoldina*, **1999**, *81*, p. 137-173.
- [6] R. Berger, M. Quack, *J. Chem. Phys.*, **2000**, *112*, p. 3148-3158.
- [7] M. Quack, J. Stohner, *Phys. Rev. Lett.*, **2000**, *84*, p. 3807-3810.
- [8] R. Berger, M. Quack, J. Stohner, *Angew. Chem.-Int. Edit.*, **2001**, *40*, p. 1667-1670.
- [9] M. Gottselig, D. Luckhaus, M. Quack, J. Stohner, M. Willeke, *Helv. Chim. Acta*, **2001**, *84*, p. 1846-1861.
- [10] R. Berger, M. Gottselig, M. Quack, M. Willeke, *Angew. Chem.-Int. Edit.*, **2001**, *40*, p. 4195-4198.
- [11] M. Quack, *Chem. Phys. Lett.*, **1986**, *132*, p. 147-153.
- [12] M. Quack, *Angew. Chem.-Int. Edit.*, **2002**, (in press), and literature cited therein.
- [13] M. Quack and J. Stohner, *Chirality*, **2001**, *13*, p. 745-753, and literature cited therein.
- [14] M. Quack, *Angew. Chem.-Int. Edit* **1989**, *28*, p. 571-586.



- [15] M. Quack, "The symmetries of time and space and their violation in chiral molecules and molecular processes", p. 172-208, in "Conceptual Tools for Understanding Nature. Proc. 2nd Int. Symp. of Science and Epistemology Seminar, Trieste April 1993", (Eds. G. Costa, G. Calucci, M. Giorgi), World Scientific Publ.: Singapore **1995**.
- [16] M. Quack, "Molecular femtosecond quantum dynamics between less than yoctoseconds and more than days: Experiment and theory", p. 781-818, in "Femtosecond Chemistry, Proc. Berlin Conf. Femtosecond Chemistry, March 1993" (Eds.: J. Manz, L. Woeste), Verlag Chemie, Weinheim, **1994**.

## ON CHARACTERIZATION OF 2-D PROTEOMICS MAPS

Milan Randić

National Institute of Chemistry, Ljubljana, Slovenia

Proteome has been defined as the totality of all proteins in cells of particular organ or tissue of living systems and proteomics research is concerned with identification, characterization, quantification and the study of interaction and function of proteins in a cell. Formally proteome is defined as PROTEin complement expressed by genOME (or tissue). The challenge is due to complexity of cells that typically contains thousands and thousands of proteins. Proteomics map is obtained by separating proteins by charge (electrophoretically) and by mass (chromatographically) on a 2-D gel. Such 2-D maps have been, and continue to be, examined mostly visually. I will review recent development of quantitative (numerical) approach for comparison of different 2-D maps based on construction of sets of map invariants, which are themselves obtained by associating with a 2-D map matrices based on embedded curves and embedded graphs (based either on partial ordering of protein spots by charge and mass, or based on clustering of spots which are at small distances). Numerical characterization of proteomics maps is viewed by some as potentially a major breakthrough in the proteomics research.

Although a characterization of 2-D proteomics maps is considering just one of many aspects of the proteomics research it is likely to help in answering other complex problems, including the study of the function of interacting proteins within a single cell. The question "Where do we come from and where are we going to?" relating to Life on a macroscale may well benefit from the question "Where do we come from and where are we going to?" relating to Cells on a microscale - which is the central question to the Proteome Research.

## **SOLAR INFLUENCE ON CLIMATE**

Henrik Svensmark  
Danish Space Research Institute

More and more studies indicate that variations in solar activity have had a significant influence on Earth's climate. In spite of many ideas and much effort, the mechanisms responsible for solar influence on climate is still not known. One possibility is to influence the atmospheric transparency by changing cloud properties via cosmic ray ionisation (the latter is modulated by solar activity). Support for this idea is found from satellite observations of cloud cover. Such data have revealed a striking correlation between the intensity of galactic cosmic rays (GCR) and low liquid clouds ( $< 3$  km). GCR is responsible for nearly all ionisation in the atmosphere below 35 km. One explanation could involve ion-induced formation of aerosol particles (0.001-1  $\mu$  in diameter) that can act as cloud condensation nuclei (CCN). A systematic variation in the properties of CCN will affect the cloud droplet distribution and thereby influence the radiative properties of clouds. If the GCR/Cloud link is confirmed, variations in galactic cosmic ray flux, caused by changes in the solar activity could influence Earth's radiation energy budget. In particular the magnetic flux carried by the solar wind has more than doubled during the last 100 years, at the same time as the global temperature has risen.

## LIPID VESICLES AND THE ORIGIN OF SOME CELLULAR PROCESSES

Saša Svetina and Boštjan Žekš

Institute of Biophysics, Faculty of Medicine, University of Ljubljana and J. Stefan  
Institute, Ljubljana, Slovenia

Lipid vesicles are macroscopic objects defined by a lipid membrane enclosing an aqueous medium and separating it from the external aqueous medium. The division of space into internal and external compartments is also the basic characteristic of living cells. Lipid vesicles can form spontaneously and as such they may have been one of the important basic structures involved in biogenesis. Lipid vesicles also possess a variety of properties which could have been employed in the further evolution of biological systems. Here it is argued that the cellular processes which involve changes of membrane conformations have their origin in the general shape behavior of closed lamellar membranes. This notion is based on some general features of the vesicle shape behavior which are the consequence of the layered membrane structure and do not depend on the structural and compositional details of the composing monolayers. One can thus ascribe these features also to prebiotic vesicles, even without knowing from what type of lipids they were made. The two general features of vesicle shape behavior that may be particularly important bases for possible physical mechanisms underlying different cellular processes are the symmetry breaking in vesicle shape transformations and the propensity of lipid vesicles for budding. The occurrence of stable asymmetric vesicle shapes is proposed to represent the mechanical origin of cellular polarity, and through this also of the biological order. Membrane budding is an obligatory step in vesicle fission and fusion processes taking place in cellular membrane trafficking. At the phenomenological level certain aspects of all these processes can be interpreted already on the basis of the understanding of the shape behavior of lipid vesicles. However, lipid vesicles and cellular systems differ in that the processes involving vesicles are in general stochastic, whereas the processes in a cell are based on a complex protein machinery and are in general deterministic. The point is made that some cellular processes, notable examples are cytokinesis, endocytosis and exocytosis, still emerged from the corresponding physical processes occurring also at the level of pure vesicles, but were properly upgraded and became well regulated in the course of evolution.

## NEW INTERSTELLAR AND LABORATORY MOLECULES

Patrick Thaddeus

Harvard-Smithsonian Center for Astrophysics 10 Garden St., Cambridge, MA, USA

Over the past five years, 130 new molecules have been discovered in our laboratory, in support of an observational program to identify and characterize new molecules in space. Most of these new species are highly reactive carbon chains -a type of molecule known to form readily and survive in the harsh, ultra-low-density environment of the interstellar gas--but a significant fraction are rings, or rings attached to carbon chains, and a number contain the hetero atoms N, O, S, and Si. On the basis of our laboratory frequencies, at least 7 of these molecules have already been identified in our galactic system in interstellar molecular clouds or circumstellar shells, and it is likely that many more will be detected during the next few years with the operation of powerful radio telescopes now under construction. The use of Fourier Transform Microwave Spectroscopy in the radio band, and laser Cavity Ringdown Spectroscopy in the optical, which yielded the new molecules will be described.

**DIFFRACTION OF MATTER WAVES FROM NANOSTRUCTURES:  
NEW PHYSICS AND NOVEL APPLICATIONS**

J. Peter Toennies  
Max-Planck-Institut für Strömungsforschung  
D-37073 Göttingen, Germany

Recent nanotechnology advances have made it possible to produce free standing transmission gratings with slits of 50 nm width. The diffraction of atoms, molecules and small clusters from such gratings provide beautiful textbook examples of matter wave phenomena [1]. Since only the wave nature of the particles is involved even the exceedingly weakly bound He dimers (binding energy  $\approx 10^{-3}$  K ( $\cong 10^{-7}$  eV)) and other small He clusters with up to 26 atoms can be separated by virtue of their different de Broglie wavelengths. Deviations of the diffraction intensities from predictions by traditional optics have been used to measure for the first time the long range particle-surface potentials [2] and the geometrical sizes of the He dimer ( $\approx 50$  Å) [3] and of the trimer (10 Å).

With three identical transmission gratings a Mach-Zehnder universal matter wave interferometer has been demonstrated. This device opens up a wide range of intriguing new experiments in molecular physics and has long range implications for studying decoherence as predicted by modern quantum theories of gravity. Transmission Fresnel zone plates have also been used to focus an atomic beam onto a 1 micron diameter spot [4], so that an essential part of a uniquely surface sensitive atom microscope is now available.

- [1] W. Schöllkopf and J.P. Toennies, *Science* **266**, 1345 (1994)
- [2] R.E. Grisenti, W. Schöllkopf, J.P. Toennies, G.C. Hegerfeldt and T. Köhler, *Phys. Rev. Lett.* **83**, 1755 (1999)
- [3] R.E. Grisenti, W. Schöllkopf, J.P. Toennies, G.C. Hegerfeldt, T. Köhler and M. Stoll, *Phys. Rev. Lett.* **85**, 2284 (2000)
- [4] R.B. Doak, R.E. Grisenti, S. Rehbein, G. Schmahl, J.P. Toennies and Ch. Wöll, *Phys. Rev. Lett.* **83**, 4229 (1999)

Brijuni Conference VIII, Brijuni, Croatia, 26.8.-31.8.2002.

**NEW SYMMETRIES FOR FULLERENES: NEW TOOLS FOR THE  
DESCRIPTION OF CARBON ONIONS AND CARBON NANOTUBES.**

R. Twarock

City University, Department of Mathematical Sciences, Northampton Square, London  
EC1V 0HB, UK

Using some simple geometric arguments it is explained how a new mathematical tool - an affine extension of noncrystallographic Coxeter groups - can be used to construct models which describe the location of atoms in fullerenes such as Carbon onions and Carbon nanotubes. We display the point sets obtained in our models and compare with existing results. Background in group theory is not expected - all concepts will be introduced and implemented on a purely geometric level.

## **CURRENT IDEAS ABOUT THE FIRST CELLS AND THEIR PRECURSOR STRUCTURES**

Peter Walde

Institut fuer Polymere, ETH, Universitaetstrasse 6, CH 8092-Zuerich, Switzerland

Nobody knows how life originated and whether it originated on Earth or somewhere else. Since today's living systems - including simple unicellular bacteria - are all very complex, it is rather difficult to understand how a living system could originate from the non-living. Although there are a number of experimental data on the synthesis of biologically relevant molecules carried out under presumably prebiotic conditions, it is difficult to imagine how the first cell(s) formed as a result of a prebiological, chemical evolution. It may well be that the ideas about the origin of the first cells do not consider all the possibilities which are given by the self-organization of amphiphilic molecules, compounds which have water-soluble as well as water-insoluble parts in their molecular structure. In addition to the obvious role of certain amphiphiles in the formation of the lipid matrix of the cell membrane, there may be other important roles of amphiphiles to be considered. Some of these possible roles are discussed.



## HIGH-PRESSURE CESIUM DISCHARGE LAMP

V. Živčec, T. Ban, H. Skenderović and G. Pichler

Institute Of Physics, Bijenička 46, P.O.Box 304, HR-10001 Zagreb, Croatia

High-pressure sodium sources of light have been spread all over the world despite the fact that color reproduction is relatively low. In addition, because of small amount of mercury in the discharge plasma, there is an environmental problem with the poisoning effect of the soil where the old lamps are stored.

In search for a better and nontoxic new light source we received pulsed high pressure cesium lamps of 80 Watt power. Color rendering index of this high-pressure cesium lamp is more than 95 (class 1A). In a usual pulsed regime the visible spectrum can be approximated with a Planck function for black body radiation at 3800 K [1,2]. Unfortunately, the efficiency of this high-pressure cesium discharge lamp has reached only 50 lumen/watt. We believe that the main cause of this might be a considerable loss in the infrared spectral region. Thus, the present application of high pressure pulsed cesium discharge lamp as a new light source is precluded.

The goal of a present research is to analyze spectra near UV, visible and infrared regions, in order to perform lamp tailoring with our knowledge of spectral phenomena in alkali plasmas [3-5]. We also believe that a few technical modifications are required to make this source of light more efficient. The new results will be presented and discussed.

### REFERENCES:

- [1] J.Liu, Ph. D. Thesis, Berlin (1998)
- [2] H.Gu, M.E.Muzeroll, J.C. Chamberlain and J.Maya, Plasma Sources Sci. Technol. 10, 1(2001)
- [3] D.Veža, R.Beuc, S.Milošević, G.Pichler, Eur. Phy. J.D 2, 45(1998)
- [4] T.Ban, H.Skenderović, S.Ter-Avetisyan and G.Pichler, Appl.Phy. B, 72, 337(2001)
- [5] R.Beuc, H.Skenderović, T.Ban, D.Veža, G.Pichler and W.Meyer, Eur.Phys. J.D 15, 209(2001)

### **3. List of Participants**

Gustaf Arrhenius  
University of California at San Diego  
9500 Gilman Drive  
La Jolla, CA 92093, USA  
email: arrhenius@ucsd.edu

Duje Bonacci  
R. Boskovic Institute  
Bijenicka 54  
Zagreb, Croatia  
email: dbonacci@eskola.hfd.hr

D. Aumiler  
Institute of Physics, Bijenička 46, P.O.  
Box 304  
HR-10001 Zagreb, Croatia  
e-mail: aumiler@ifs.hr

S. Danko Bosanac  
R. Boskovic Institute  
Bijenicka 54  
Zagreb, Croatia  
email: danko.bosanac@irb.hr

Jadson C. Belchior  
Universidade Federal de Minas Gerais  
Av. Antonio Carlos 6627  
Departamento de Quimica - ICEx  
31.270-901 - Belo Horizonte  
Minas Gerais, BRASIL  
email: jadson@carbono.qui.ufmg.br

Charles Cantor  
Sequenom Inc.  
3595 John Hopkins Court  
San Diego, CA 92121-1331  
USA  
email: ccantor@sequenom.com

Robert Beuc  
Institute of Physics, Bijenička 46, P.O.  
Box 304  
HR-10001 Zagreb, Croatia  
email: beuc@ifs.hr

Ludwik M. Celnikier  
Observatoire de Meudon  
92195 Meudon  
France  
email: ludwik.celnikier@obspm.fr

A.E. Blaugrund  
Department of Particle Physics  
Weizmann Institute of Science  
IL76100 Rehovot  
ISRAEL  
email: fnbla@weizmann.ac.il

Vladimir Damgov  
The Bulgarian Academy of Sciences  
Space Research Institute  
6 Moscovska St., POBox 799  
1000 Sofia, Bulgary  
email: vdamgov@bas.bg

Leo Blitz  
University of California at Berkeley  
Radio Astronomy Laboratory  
635 Campbell 3411  
Berkeley, CA-94720-3411  
USA  
email: blitz@gmc.Berkeley.edu

Pier Paolo Delsanto  
Dipartimento di Fisica  
Politecnico di Torino  
C. Duca Abruzzi 24  
Torino  
10129 ITALY  
email: delsanto@polito.it

M.P.Dos Santos  
Physics Department  
University of Évora  
Colegio Luís A. Verney  
7000-671 ÉVORA, PORTUGAL  
email: mpsantos@uevora.pt

Martin Harwit  
Cornell University  
Department of Astronomy  
Ithaca  
NY 14853-6801, USA  
email: harwit@bellatlantic.net

Nadja Došlić  
R. Bošković Institute  
Bijenička 54  
Zagreb, Croatia  
email: nadja.doslic@irb.hr

Saša Ilijić  
Geodetski fakultet, University of  
Zagreb  
Kačićeva 26  
Zagreb, Croatia  
email: silijic@geof.hr

Jean Durup  
IRSAMC UPS  
118 rte de Narbonne  
31062 Toulouse Cedex, France  
email: durup@irsamc.ups-tlse.fr

Nsiah Yeboah Isaac  
RM 375 Unity Hall U S T,  
Kumasi A/R  
Ghana W/A  
email: destiny1555a@yahoo.com

Thomas Ebbesen  
Laboratoire des Nanostructures  
Université Louis Pasteur  
4 rue B. Pascal  
67000 Strasbourg, France  
email: twe@chimie.u-strasbg.fr

Michael Jura  
Department of Physics and Astronomy  
University of California  
Box 951547  
Los Angeles, CA 90095-1547, USA  
email: jura@astro.ucla.edu

Manfred Eigen  
MPI für Biophysikalische Chemie  
37070 Göttingen  
Germany  
email: manfred.eigen@gwdg.de

Davor Juretić  
Department of Physics, University of  
Split  
Nikole Tesle 12, Split, Croatia  
email: juretic@pmfst.hr

Professor Enzo Gallori  
Department of Animal Biology and  
Genetics,  
University of Florence  
Via Romana 17/19  
50125 Florence, Italy  
email: gallori@dbag.unifi.it

Dubravko Klabučar  
Department of Physics  
University of Zagreb  
Bijenicka 32  
Croatia  
email: klabucar@sirius.phy.hr

Ante Graovac  
R. Boskovic Institute  
Bijenicka 54  
Zagreb, Croatia  
email: graovac@rudjer.irb.hr

Leo Klasinc  
R. Boskovic Institute  
Bijenicka 54  
Zagreb, Croatia  
email: klasinc@joker.irb.hr

Bill Klemperer  
Harvard University  
Department of Chemistry  
12 Oxford Street, Cambridge  
Mass. 02138, USA  
email: klemperer@chemistry.harvard.edu

Goran Kovačević  
R. Bošković Institute  
Bijenička 54  
Zagreb, Croatia  
email: gkova@irb.hr

Tomislav Krčmar  
Croatian Radio Television  
Prisavlje 3  
10 000 Zagreb, Croatia

Harry Kroto  
University of Sussex  
Falmer, Brighton  
BN1 9RH, UK  
email: kroto@sussex.ac.uk

Krešimir Kumerički  
Department of Physics  
University of Zagreb  
Bijenička 32, Zagreb, Croatia  
email: kkumer@phy.hr

Sydney Leach  
Laboratoire Photophysical Molecular  
Universite de Paris-Sud, Bat 213  
F-91405 Orsay, France  
email: sydney.leach@obspm.fr

Jean - Marie Lehn  
ISIS-ULP, UMR CNRS 7006  
4 rue Blaise Pascal  
F-67000 Strasbourg, France  
email: lehn@chimie.u-strasbg.fr

Ivan Ljubić  
R. Bošković Institute  
Bijenička 54  
Zagreb, Croatia  
email: iljubic@rudjer.irb.hr

Joao Magueijo  
Imperial College of Science,  
Technology and Medicine  
Department of Physics  
Exhibition Road  
London SW7 2AZ, UK  
email: j.magueijo@ic.ac.uk

Klaus Mainzer  
Universität Augsburg  
Universität Str. 10  
D-86159 Augsburg, Germany  
email: klaus.mainzer@phil.uni-augsburg.de

Mirjana Maksić  
R. Bošković Institute  
Bijenička 54  
Zagreb, Croatia  
email: mmaksic@emma.irb.hr

Zvonimir B. Maksić  
R. Bošković Institute  
Bijenička 54  
Zagreb, Croatia  
email: zmaksic@spider.irb.hr

Mladen Martinis  
R. Bošković Institute  
Bijenička 54  
Zagreb, Croatia  
email: martinis@thphys.irb.hr

Ivana Matanović  
Chemistry Department  
University of Zagreb  
Zagreb, Croatia  
email: ivana@alba.irb.hr

Michel Mayor  
Observatoire de Geneve  
Université de Geneve  
CH-1290 Versoix  
Switzerland  
email: Michel.Mayor@obs.unige.ch

Ivica Picek  
Odsjek teorijske fizike  
Bijenička c. 32  
POB 162  
10 000 Zagreb  
email: picek@phy.hr

Vesna Mikuta-Martinis  
R. Bošković Institute  
Bijenička 54  
Zagreb, Croatia  
email address: mikutama@thphys.irb.hr

Goran Pichler  
Institute of Physics  
Bijenička 46, P.O. Box 304  
HR-10001 Zagreb, Croatia  
email: pichler@ifs.hr

Mira Morović  
Institut za oceanografiju i ribarstvo  
P.P. 500  
21000 Split, Hrvatska  
email: morovic@izor.hr

Teuta Pilzota  
The Clarendon Laboratory  
Condensed Matter Physics  
University of Oxford  
Parks Road, Oxford OX1 3PU, UK

Mladen Movre  
Institute of Physics  
Bijenička 46, P.O. Box 304  
HR-10001 Zagreb, Croatia  
email: movre@ifs.hr

Levon Pogossian  
Blackett Laboratory  
Imperial College  
Prince Consort Road  
London, SW7 2BW, U.K.  
email: l.pogossian@ic.ac.uk

John N. Murrell  
University of Sussex  
Falmer, Brighton  
BN1 9RH, UK  
email: j.n.murrell@sussex.ac.uk

Evgeniy D. Prudnikov  
Deptment of Physics  
University of St-Petersburg  
Universitetskaya 7/9, 199034, Russia  
email: evgeniy@EP2256.spb.edu

Sonja Nikolić  
R. Bošković Institute  
Bijenička 54  
Zagreb, Croatia  
email: sonja@rudjer.irb.hr

Martin Quack  
Laboratorium für Physikalische  
Chemie, ETH Zürich Zentrum  
CH-Zürich, Switzerland  
email: martin@quack.ch

Vesna Noethig-Laslo  
R. Bošković Institute  
Bijenička 54  
Zagreb, Croatia  
email: laslo@faust.irb.hr

Milan Randić  
National Institute of Chemistry  
Ljubljana, Slovenia  
email: milan.randic@KI.si

Alexandar Sabljic  
R. Bošković Institute  
Bijenička 54  
Zagreb, Croatia  
sablji@rudjer.irb.hr

Nenad Trinajstić  
R. Bošković Institute  
Bijenička 54  
Zagreb, Croatia  
email: trina@rudjer.irb.hr

Flavio Scappini  
ISM - CNR  
Via P. Gobetti, 101  
40129 Bologna, Italy  
email: F.Scappini@ism.bo.cnr.it

Reidun Twarock  
City University  
Department of Mathematical Sciences  
Northampton Square  
London EC1V 0HB, UK  
email: r.twarock@city.ac.uk

Giacinto Scoles  
Chemistry Department and Princeton  
Materials Institute, Princeton  
University  
Princeton, New Jersey 08544, USA  
email: gscoles@Princeton.EDU

Dejan Vinković  
Department of Physics and Astronomy  
University of Kentucky  
177 CP-bldg  
Lexington, KY 40506-0055  
e-mail: dejan@pa.uky.edu

Henrik Svensmark  
Danish Space Research Institute  
Juliane Maries Vej 30  
DK-2100 Copenhagen, Denmark  
email: hsv@dsri.dk

Ivana Vlah  
R. Bošković Institute  
Bijenička 54  
Zagreb, Croatia  
email: ivlah@irb.hr

Saša Svetina  
Institute of Biophysics  
Faculty of Medicine, University of  
Ljubljana  
Lipičeva 2, 1000 Ljubljana, Slovenia  
email: sasa.svetina@biofiz.mf.uni-lj.si

Peter Walde  
Institut fuer Polymere  
ETH-Zentrum  
Univerisität Strasse 6  
CH-8092 Zürich, Switzerland  
email: pwalde@ifp.mat.ethz.ch

Patrick Thaddeus  
Harvard-Smithsonian Center for  
Astrophysics  
10 Garden St., Cambridge, MA, USA  
email: pthaddeus@cfa.harvard.edu

Ruthild Winkler-Owatitsch  
MPI für Biophysikalische Chemie  
37070 Göttingen  
Germany

Jan Peter Toennies  
MPI für Strömungsforschung  
Bunsenstrasse 10  
D-37073 Göttingen, Germany  
email: jtoenni@gwdg.de

Ingrid Wysong  
European Office of Aerospace  
Research and Development  
223/231 Old Marylebone Rd.  
London, NW1 5TH, UK  
email: ingrid.wysong@london.af.mil

V. Živčec  
Institute of Physics  
Bijenička 46, P.O. Box 304  
HR-10001 Zagreb, Croatia  
e-mail: aumiler@ifs.hr

Tomislav Živković  
R. Bošković Institute  
Bijenička 54  
Zagreb, Croatia  
email: zivkovic@rudjer.irb.hr



**ISBN 953-6690-23-3**